

Chaotic dynamics and superdiffusion in a Hamiltonian system with many degrees of freedom

V. Latora¹ and A. Rapisarda²

Dipartimento di Fisica Università di Catania, Corso Italia 57 I 95129 Catania, Italy

S. Ruffo³

*Laboratoire de Physique, Ecole Normale Supérieure de Lyon
46 allée d'Italie 69364 Lyon Cedex 07 France, INFN and INFN Firenze, Italy*

Abstract

We discuss recent results obtained for the Hamiltonian Mean Field model. The model describes a system of N fully-coupled particles in one dimension and shows a second-order phase transition from a clustered phase to a homogeneous one when the energy is increased. Strong chaos is found in correspondence to the critical point on top of a weak chaotic regime which characterizes the motion at low energies. For a small region around the critical point, we find anomalous (enhanced) diffusion and Lévy walks in a transient temporal regime before the system relaxes to equilibrium.

Key words: Hamiltonian dynamics, deterministic chaos, Lyapunov exponents, relaxation to equilibrium, anomalous diffusion, Lévy walks

PACS numbers: 05.45.Pq, 05.20.-y, 05.60.cd, 05.70.Fh

1 Introduction

In this contribution we present a novel study of chaos and anomalous diffusion in the dynamics of the single-particle motion in the Hamiltonian Mean Field

¹ E-mail: latora@ct.infn.it

² E-mail: andrea.rapisarda@ct.infn.it

³ On leave from Dipartimento di Energetica “S. Stecco”, Università di Firenze, Via S. Marta, 3 I-50139, Firenze, Italy, E-mail: ruffo@avanzi.de.unifi.it

(HMF) model. This model describes N classical particles moving on the unit circle and interacting through an infinite range potential. It has been recently studied both analytically and numerically [1–4]. At low energy the particles are clustered and, as the energy is increased, the system undergoes a second-order phase transition, reaching a high temperature “gaseous” phase where the particles are homogeneously distributed on the circle. The motion is chaotic and the Lyapunov exponent reaches a maximum at the critical point.

Particle motion is superdiffusive for a range of energies close to the critical point, but only in a transient quasi-stationary regime preceding equilibration. Particles can evaporate from the cluster showing Lévy walks. Close to the critical point, if the continuum ($N \rightarrow \infty$) limit is performed before the $t \rightarrow \infty$ limit, canonical equilibrium is never reached and quasi-stationary superdiffusive states live forever [1–4]. Though anomalous diffusion and Lévy walks have been widely studied in the recent literature in models with a few degrees of freedom [5,6], there are very few studies for many-degrees-of-freedom systems [7,8]. The HMF model has the advantage of having an exact solution in the canonical ensemble, and therefore microscopic dynamics can be put in connection to macroscopic features, thus offering a very interesting perspective [3,4].

In the following sections we remind the reader the details of the model and then we discuss superdiffusion in connection to microscopic chaotic dynamics.

2 The model

HMF describes a system of N classical particles moving on a circle, characterized by the angles θ_i and the conjugate momenta p_i . Each particle interact with all the others according to the following Hamiltonian:

$$H(\theta, p) = K + V, \quad (1)$$

where

$$K = \sum_{i=1}^N \frac{p_i^2}{2} \quad V = \frac{1}{2N} \sum_{i,j=1}^N [1 - \cos(\theta_i - \theta_j)] \quad (2)$$

are the kinetic and potential energy. If we define a spin vector associated to each particle $\mathbf{m}_i = [\cos(\theta_i), \sin(\theta_i)]$ and a total magnetization $\mathbf{M} = \frac{1}{N} \sum_{i=1}^N \mathbf{m}_i$ the Hamiltonian then describes N classical fully coupled spins, similarly to the XY model. With this interpretation, the system is a ferromagnet at low




Fig. 1. Total magnetization M as a function of the energy per particle $U = E/N$. Numerical simulations are compared with the canonical prediction (full curve). The insets show a schematic pictorial view of the condensed and the homogeneous phase. See text.

energy and shows a second-order phase transition at the critical energy $U_c = E_c/N = 0.75$ (corresponding to the critical temperature $T_c = 0.5$ in the canonical ensemble) [1–3]).

We show in fig.1 the total magnetization vs. energy $U = E/N$. Numerical simulations (dots) are compared with the canonical ensemble result (full line). Pictorial views of the low energy ferromagnetic (clustered) phase and high energy paramagnetic (gaseous) phases are shown in the insets. The equations of motion for the N particles are

$$\dot{\theta}_i = p_i, \quad \dot{p}_i = -M \sin(\theta_i - \phi) \quad , \quad i = 1, \dots, N \quad , \quad (3)$$

where (M, ϕ) are respectively the modulus and the phase of the total magnetization vector $\mathbf{M}$. These equations are formally equivalent to those of a perturbed pendulum. To study relaxation to canonical equilibrium, we solve these equations on the computer using fourth order symplectic algorithms (the details can be found in Refs [2,3]). We start the system in a given initial state and we compute θ_i, p_i at each time step, and from them the total magnetization M and temperature T (through the relation $T = 2 \langle K \rangle / N$). We consider systems with an increasing size N and different energies $U = E/N$. The single-particle motion is given by Eq. (3), if we consider the mean-field pair (M, ϕ) as an *external* time dependent forcing. In the next section we focus our attention on the dynamics of the single-particle motion and we show that




Fig. 2. Single-particle phase space for three different energy regimes: $U = 0.1$ ($< U_c$), $U = 0.69$ (close to U_c) and $U = 1$ ($> U_c$). See text for more details.

it is chaotic and superdiffusive.

3 Chaotic motion and superdiffusion

In this section we discuss the numerical results. In fig. 2 we show single-particle phase spaces for three different energies: below (2a), close to (2b) and above (2c) the critical energy $U_c = 0.75$. While for $U = 0.1$ the particle never escapes the cluster and moves on the chaotic web near the center of the nonlinear resonance, close to the critical point (at $U_c = 0.69$) it escapes frequently from the cluster and is then trapped again, exhibiting the well known phenomenon of *separatrix crossing*. Above the critical energy one has mainly a ballistic regime, separatrix crossings are rarely observed and particles interact very weakly.

We show in fig. 3 the largest Lyapunov exponent λ_{sp} for the single-particle motion as a function of the total energy per particle. The motion is chaotic and has a peak at the critical point, but λ_{sp} is much smaller than the largest Lyapunov exponent calculated for the full dynamics in Refs. [2,3]. The reason




Fig. 3. (a) We show the largest Lyapunov exponent for the single-particle as a function of U . In the inset we show the behavior at small energy in a log-log plot, see text. (b) We report numerical simulations (open circles) that show that λ_{sp} scales like $N^{-1/3}$ (straight line).

is that in the calculation of the Lyapunov of the full system one gets the additional contribution to the instability of the off-diagonal terms of the Jacobian. We show in the inset a log-log plot with a fit at energies less than 0.3, which is a power law, i.e. $\lambda \sim U^{1/3}$; again quite far from the $\sqrt{U}$ law found for the full dynamics. A very weak N dependence is present in this region. In the high energy region λ_{sp} vanishes with N , with the “universal” law $\lambda_{sp} \sim N^{-1/3}$, which is found in many different contexts [9]. This result is displayed in fig.3(b) for $U = 2$.

Diffusion and transport of a particle in a medium or in a fluid flow are characterized by the mean square displacement $\sigma^2(t)$ in the long-time limit. In general, one has

$$\sigma^2(t) \sim t^\alpha \quad (4)$$

with $\alpha = 1$ for normal diffusion. All the processes with $\alpha \neq 1$ are termed anomalous diffusion, namely subdiffusion for $0 < \alpha < 1$ and superdiffusion for $1 < \alpha < 2$.

In order to study anomalous diffusion in HMF we follow the dynamics of each particle starting the system in a “water bag”, i.e. a far-off-equilibrium initial condition obtained by putting all the particles at $\theta_i = 0$ and giving them a uniform distribution of momenta with a finite width centered around zero. We compute the variance of the one-particle angle θ according to the expression

$$\sigma^2(t) = \langle (\theta - \langle \theta \rangle)^2 \rangle \quad , \quad (5)$$




Fig. 4. (a) Lévy walks for different test particles at $U = 0.69$. (b) Variance of the single-particle displacement as a function of time for four different energies.

where $\langle \cdot \rangle$ indicates the average over the N particles, and we fit the value of the exponent α in Eq. (4).

In fig. 4(a) we show the motion of four particles in the same simulation as a function of time for $N = 500$ and $U = 0.69$. The figure shows that the erratic motion is not brownian and Lévy walks are present. Each particle experiences in time localized motion and free walks. In fig.4(b) we show the mean square displacement as a function of time. For energies close to the critical one the exponent α differs from 1, as it should be for normal diffusion, and takes values close to 1.4. One has therefore superdiffusion, which however turns again into normal after a crossover time. This behavior does not seem to depend strongly on energy, although the superdiffusion exponent α is slightly energy dependent. In Ref. [4] we have shown that this crossover time coincides with the equilibration one and that by using the model of Ref. [6] it is possible to interpret this dynamics in terms of Lévy walks.

4 Conclusion

We have discussed chaotic and superdiffusive motion in a N -body Hamiltonian system which exhibits a second-order phase transition. The single-particle motion is shown to be chaotic for energies smaller than the critical one. Superdiffusion is present in a transient out-of-equilibrium regime for energies close to the critical one, where the system is strongly chaotic and quasi-stationary states exist. The crossover time from enhanced diffusion to normal coincides with the equilibration time which diverges with N . This is true for fully coupled systems, but should appear also if the interaction decays with the dis-

tance among the particles with a power law [10]. Our model lies in the class of self-consistent many-degrees-of-freedom dynamics, to which also belongs the single-wave model described in Ref. [11]. This is the closest contact of our results to the Lévy-walk behavior found in fluid experiments [12].

Acknowledgements

We dedicate this work to the memory of Giovanni Paladin who was an enthusiastic pioneer in the study of chaos and statistical mechanics, and the first one to explain to one of the authors (S.R.) the concept of anomalous diffusion. We thank C. Anteneodo, M. Antoni, M. Baranger, E. Barkai, D. del-Castillo-Negrete, P. Grigolini, A. Torcini and C. Tsallis for useful discussions. S.R. thanks ENS-Lyon for financial support.

References

- [1] S. Ruffo, *Transport and Plasma Physics*, S. Benkadda, Y. Elskens and F. Doveil eds., World Scientific, Singapore (1994), p. 114; M. Antoni and S. Ruffo, *Phys. Rev. E* 52 (1995) 2361.
- [2] V. Latora, A. Rapisarda and S. Ruffo, *Phys. Rev. Lett.* 80 (1998) 692.
- [3] V. Latora, A. Rapisarda and S. Ruffo, *Physica D* 131 (1999) 38.
- [4] V. Latora, A. Rapisarda and S. Ruffo, *Phys. Rev. Lett.* 83 (1999) 2104.
- [5] M.F. Shlesinger, G.M. Zaslavsky and U. Frisch Eds., *Lévy flights and related topics*, Springer-Verlag Berlin (1995) and Refs. therein.
- [6] J. Klafter and G. Zumofen, *Phys. Rev. E* 49 (1994) 4873.
- [7] K. Kaneko and T. Konishi, *Phys. Rev. A* 40, (1989) 6130; T. Konishi, *Progr. Theor. Phys.* 98, (1989) 19.
- [8] M. Antoni and A. Torcini, *Phys. Rev. E* 57, R6233 (1998) R6233; A. Torcini and M. Antoni, *Phys. Rev. E* 59, (1999) 2746.
- [9] M. Antoni, S. Ruffo and A. Torcini, to be published in *The Chaotic Universe*, V. Gutzwiller ed., World Scientific (2000)
- [10] C. Anteneodo and C. Tsallis, *Phys. Rev. Lett.* 80 (1998) 5313.
- [11] D. del-Castillo-Negrete, *Phys. Lett. A* 241 (1998) 99; *Physics of Plasmas* 5 (1998) 3886.
- [12] T.H. Solomon, E.R. Weeks and H.L. Swinney, *Phys. Rev. Lett.* 71 (1993) 3975.